

REVIEW OF MOBIUS TRANSFORMATION

KarthikVeduar ¹, YashShirdhankar ², SwapnilYadav ³, Ashutosh
Bhosale ⁴, SakibMalim ⁵
¹⁻⁵ Students

Department of Information Technology
K.C. College of Engineering and Management Studies and Research, Thane
Mumbai, India

Abstract:

Conformal (Same form or shape) mapping is an important technique used in complex analysis and has many applications in different physical situations. If the function is harmonic (ie it satisfies Laplace's equation $\nabla^2 f = 0$) then the transformation of such functions via conformal mapping is also harmonic. So equations pertaining to any field that can be represented by a potential function (all conservative fields) can be solved via conformal mapping. If the physical problem can be represented by complex functions but the geometric structure becomes inconvenient then by an appropriate mapping it can be transferred to a problem with much more convenient geometry. This article gives a brief introduction to conformal mappings and some of its applications in physical problems.

I. INTRODUCTION

A conformal map is a function which preserves the angles. Conformal map preserves both angles and shape of infinitesimal small figures but not necessarily their size. More formally, a map

$$w = f(z) \quad \text{--(1)}$$

is called conformal (or angle-preserving) at z_0 if it preserves oriented angles between curves through z_0 , as well as their orientation, i.e., direction.

An important family of examples of conformal maps comes from complex analysis. If U is an open subset of the complex plane, then a function

$$f: U \rightarrow \mathbb{C}$$

is conformal if and only if it is holomorphic and its derivative is everywhere non-zero on U . If f is antiholomorphic (that is, the conjugate to a holomorphic function), it still preserves angles, but it reverses their orientation. The Riemann mapping theorem, states that any non- empty open simply connected proper subset of \mathbb{C} admits a bijective conformal map to the open unit disk (the open unit disk around P (where P is a given point in the plane), is the set of points whose distance from P is less than 1)




FIG. 1: Mapping of graph

in complex plane C i.e., if U is a simply connected open subset in complex plane C , which is not all of C , then there exists a bijective i.e., one-to-one mapping f from U to open unit disk D .

$$f: U \rightarrow D \text{ where} \\ D = \{z \in C: |z| < 1\}$$

As f is a bijective map it is conformal.

A map of the extended complex plane (Which is conformally equivalent to a sphere) onto itself is conformal if and only if it is a Mobius transformation i.e., a transformation leading to a rational function of the form $f(z) = \frac{az+b}{cz+d}$. Again, for the conjugate, angles are preserved, but orientation is reversed.

II. BASIC THEORY

Let us consider a function

$$w = f(z) \quad \text{---} \quad (2)$$

where $z = x + iy$ and $w = u + iv$ We find that
 $dz = dx + idy$ and $dw = du + idv$

$$|dz|^2 = dx^2 + dy^2, \text{-----} (3)$$

and

$$|dw|^2 = du^2 + dv^2 \text{-----} (4)$$

Then the square of the length element in (x,y) plane is

$$ds^2 = dx^2 + dy^2 \text{-----} (5)$$

and square of length element in (u,v) plane is

$$ds^2 = du^2 + dv^2 \text{-----} (6)$$

From equations (3), (4), (5), (6) we find that,

$$dS/ds = |dw/dz| \text{-----} (7)$$

ie the ratio of arclengths of two planes remains essentially constant in the neighborhood of each point in z plane provided $w(z)$ is analytic and have a nonzero or finite slope at that point. This implies the linear dimensions in two planes are proportional and the net result of this transformation is to change the dimensions in equal proportions and rotate each infinitesimal area in the neighborhood of the point. Thus the angle (which is represented as the ratio of linear dimensions) is preserved although shape in a large scale will not be preserved in general as the value of $|dw/dz|$ will vary considerably at different

points in the plane. Due to this property such transformations are called conformal. This leads to the following theorem.

Theorem : Assume that $f(z)$ is analytic and not constant in a domain D of the complex z plane. For any point $z \in D$ for which $f'(z) \neq 0$, this mapping is conformal, that is, it preserves the angle between two differentiable arcs.

Example: Let D be the rectangular region in the z plane bounded by $x=0, y=0, x=2$ and $y=1$. The image of D under the transformation $w = (1+i)z + (1+2i)$ is given by the rectangular region D' of the w plane bounded by $u+v=3, u-v=-1, u+v=7$ and $u-v=-3$.

If $w = u+iv$, where $u, v \in \mathbb{R}$, then $u = x - y + 1, v = x + y + 2$. Thus the points $a, b, c,$ and d are mapped to the points $(0,3), (1,2), (3,4),$ and $(2,5)$, respectively. The line $x=0$ is mapped to $u = -y + 1, v = y + 2$, or $u + v = 3$; similarly for the other sides of the rectangle (fig 2). The rectangle D' is translated by $(1+2i)$, rotated by an angle $\pi/4$ in the counterclockwise direction, and dilated by a factor $\sqrt{2}$. In general, a linear transformation $f(z) = \alpha z + \beta$, translates by β , rotates by $\arg(\alpha)$, and dilates (or contracts) by $|\alpha|$. Because $f'(z) = \alpha \neq 0$, a linear transformation is always conformal.




FIG. 2: Mapping of a rectangle

III . APPLICATIONS

A large number of problems arising in fluid mechanics, electrostatics, heat conduction, and many other physical situations can be mathematically formulated in terms of Laplace's equation. i.e, all these physical problems reduce to solving the equation

$$\Phi_{xx} + \Phi_{yy} = 0 \text{ ----- (8)}$$

in a certain region D of the z plane. The function $\Phi(x, y)$, in addition to satisfying this equation also satisfies certain boundary conditions on the boundary C of the region D . From the theory of analytic functions we know that the real and the imaginary parts of an analytic function satisfy Laplace's equation. It follows that solving the above problem reduces to finding a function that is analytic in D and that satisfies certain boundary conditions on C . It turns out that the solution of this problem can be greatly simplified if the region D is either the upper half of the z plane or the unit disk.

Example: Consider two infinite parallel flat plates, separated by a distance d and maintained at zero potential. A line of charge q per unit length is located between the two planes at a distance 'a' from the lower plate. The problem is to find the electrostatic potential in the shaded region of the z plane.

The conformal mapping $w = \exp(\pi z/d)$ maps the shaded strip of the z plane onto the upper half of the w plane. So the point $z = ia$ is mapped to the point $w = \exp(i\pi a/d)$; the points on the lower plate, $z = x$, and on the upper plate, $z = x + id$, map to the real axis $w = u$ for $u > 0$ and $u < 0$, respectively.

$$\Omega(w) = -2 \log(w - w_0) + 2q \log(w - \bar{w}_0) = 2q \log\left(\frac{w - \bar{w}_0}{w - w_0}\right) \tag{9}$$

Let us consider a line of charge q at w_0 and a line of charge $-q$ at \bar{w}_0 . Consider the associated complex potential




FIG. 3: Mapping of two infinite parallel conducting plate with a charge in between

Calling C_q a closed contour around the charge q , we see that Gauss law is satisfied,

$$\oint_{C_q} E_n ds = \text{Im} \oint_{C_q} \bar{E} dz = \text{Im} \oint_{\tilde{C}_q} -\Omega'(w) = 4\pi q \tag{10}$$

where \tilde{C}_q is the image of C_q in the w - plane. Then, calling $\Omega = \Phi + i\Psi$, we see that Φ is zero on the real axis of the w plane.

Consequently, we have satisfied the boundary condition $\Phi = 0$ on the plates, and hence the electrostatic potential at any point of the shaded region of the z plane is given by

$$\Phi = 2q \log\left(\frac{w - e^{-iv}}{w - e^{iv}}\right) \tag{11}$$

where $v = \pi a/d$


Conformal mappings are invaluable for solving problems in engineering and physics that can be expressed in terms of functions of a complex variable, but that exhibit inconvenient geometries. By choosing an appropriate mapping, the analyst can transform the inconvenient geometry into a much more convenient one

Conformal mapping has various applications in the field of medical physics. For example, conformal mapping is applied to brain surface mapping. This is based on the fact that any genus zero (The genus of a connected, orientable surface is an integer representing the maximum number of cuttings along closed simple curves without rendering the resultant manifold disconnected; a sphere, disk or annulus have genus zero) surface can be mapped conformally onto the sphere and any local portion thereof onto a disk.

Conformal mapping can be used in scattering and diffraction problems. For scattering and diffraction problems of plane electromagnetic waves, the mathematical problem involves finding a solution to a scalar wave function which satisfies both boundary conditions and radiation conditions at infinity. Exact solutions are available for such problems only for a few cases.

IV. CONCLUSION

There are different aspects of conformal mapping that can be used for practical applications though the essence remains the same: it preserves the angle and shape locally and mappings of harmonic potentials remains harmonic.



These properties of conformal mapping make it advantageous in complex situations, specifically electromagnetic potential problems for general systems. Various conformal techniques such as genus zero conformal mapping is also used to complex surface mapping problems. However, the conformal mapping approach is limited to problems that can be reduced to two dimensions and to problems with high degrees of symmetry. It is often impossible to apply this technique when the symmetry is broken.

REFERENCES:

- [1] Ahlfors, L., *Complex Analysis*, McGraw–Hill, New York, 1966.
- [2] Apostol, T.M., *Calculus*, Blaisdell Publishing Co., Waltham, Mass., 1967–69.
- [3] Batchelor, G.K., *An Introduction to Fluid Dynamics*, Cambridge University Press, Cambridge, 1967.
- [4] Birkhoff, G., and Rota, G.–C., *Ordinary Differential Equations*, Blaisdell Publ. Co., Waltham, Mass., 1962.
- [5] Boyce, W.E., and DiPrima, R.C., *Elementary Differential Equations and Boundary Value Problems*, 7th ed., John Wiley & Sons, Inc., New York, 2001.
- [6] Bronstein, M., *Symbolic integration I: Transcendental Functions*, Springer–Verlag, New York, 1997.
- [7] Delillo, T.K., The accuracy of numerical conformal mapping methods: a survey of examples and results, *SIAM J. Numer. Anal.* 31 (1994), 788–812.
- [8] Fine, B., and Rosenberger, G., *The Fundamental Theorem of Algebra*, Undergraduate Texts in Mathematics, Springer–Verlag, New York, 1997.
- [9] Franklin, J., Green's functions for Neumann boundary conditions, preprint, 2012, arXiv:1201.6059.
- [10] Gradshteyn, I.S., and Ryzhik, I.W., *Table of Integrals, Series and Products*, Academic Press, New York, 1965.
- [11] Hale, J.K., *Ordinary Differential Equations*, Second Edition, R.E. Krieger Pub. Co., Huntington, N.Y., 1980.
- [12] Henrici, P., *Applied and Computational Complex Analysis*, vol. 1, J. Wiley & Sons, New York, 1974.
- [13] Hutter, K., and Wang, Y., *Fluid and Thermodynamics. Volume 1: Basic Fluid Mechanics*, Advances in Geophysical and Environmental Mechanics and Mathematics, Springer International Publ., Switzerland, 2016.